



**Technique to explore
the internal structure
of an erupting
volcano**

T. Kusagaya and
H. K. M. Tanaka

Development of the very long range muographic imaging technique to explore the internal structure of an erupting volcano, Shinmoe-dake, Japan

T. Kusagaya and H. K. M. Tanaka

Earthquake Research Institute, The University of Tokyo, 1-1-1 Yayoi, Bunkyo,
113-0032 Tokyo, Japan

Received: 27 April 2015 – Accepted: 11 July 2015 – Published: 29 July 2015

Correspondence to: T. Kusagaya (kusagaya@eri.u-tokyo.ac.jp)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

Muography offers us a tool to observe hazardous erupting volcanoes remotely. However, practical muographic observations of volcanoes from a distance are difficult; therefore, various observations have been performed in the vicinity (< 1.5 km) of volcano peaks to suppress background noise and enhance images. In this study, we created a muographic image directly beneath the caldera floor of the erupting Shinmoe-dake volcano in Japan by locating our muography telescope 5 km from the peak. Shinmoe-dake volcano began to erupt on 19 January 2011, and in less than one month, the ejected lava almost completely filled the caldera and completely changed the topography of the caldera floor. The resultant image shows a low-density region underneath the western part of the newly created caldera floor, which indicates the existence of a void there. After the volcano became less active in February 2011, infrequent eruptions might have left a void beneath the caldera floor, which may trigger a collapse in the future. We anticipate that our novel muography will be a practical tool for monitoring and predicting eruption sequences in the near future.

1 Introduction

To date, one of the keys to successful volcano muography to directly contribute to the understanding the eruption dynamics has been to shorten the time required for capturing a practical radiographic image of an erupting volcano (Tanaka, 2014). For example, Tanaka et al. (2014) reduced electromagnetic background events by adding redundant detectors and radiation shields to a conventional muography telescope (e.g. Tanaka and Yokoyama, 2013), thereby improving the time resolution. They successfully visualized magma movements with a time resolution of 3 days. The reduction of background events is known to be essential for shortening the measurement time (Tanaka and Yokoyama, 2013).

GID

5, 207–246, 2015

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Another important benefit of muography is to allow us to observe hazardous volcanoes at a long distance. However, the telescopes have needed to be placed in the vicinity (0.5–1.6 km) of the volcano craters (Tanaka et al., 2014, 2010, 2009a,b, 2007a, b; De Lellis et al., 2014; Tanaka and Yokoyama, 2013; Carbone et al., 2013; Kusagaya et al., 2013; Lesparre et al., 2012; Hernández et al., 2013; Miyamoto et al., 2012) (Fig. 1). Although Cârloganu et al. (2013) located their telescope at a distance of 2 km from the peak of the Puy de Dôme volcano, they reported that the background noise dominated the muon flux when the rock thickness exceeded 1 km. Such short-range muography is practical when the target volcanoes are dormant or less active.

Thus far, there are two examples of muography measurements that targeted erupting volcanoes. In one of the examples, the telescope had been buried underground prior to the eruption (Tanaka et al., 2009b), and in the second example, the eruption was non-explosive (Tanaka et al., 2014). In the latter case, the volcanic eruption index (VEI) was zero.

In this work, we investigated possibilities of using very long-range muography (VLRM) as an alternative remote sensing tool for volcano monitoring. In conjunction with a newly developed analysis method, we used a low background muography telescope to image the density distribution beneath the caldera floor of Shinmoe-dake Volcano, Japan.

The latest eruption of the Shinmoe-dake volcano took place on 19 January 2011. An eruption column reached 3000 m a.s.l. on 26 January. A small lava dome (10 m in diameter) was created on 28 January and continued to grow inside the caldera. On 30 January, it almost completely filled the caldera. Miyazaki Prefecture reported that 1150 people had been evacuated from the area on the same day. At the same time, the area within 3 km of the mountain was designated as a forbidden zone. On 2 February, a volcanic bomb reached beyond this restriction line. It reached 3.2 km from the peak of the mountain (Japan Meteorological Agency, 2013). The amount of magma discharged between 19 January and 2 February was estimated to be $2.7\text{--}3.7 \times 10^6 \text{ m}^3$ (VEI 2).

Since 2011, the Japan Meteorological Agency (JMA) volcano eruption levels remained high and were Level 2 in 2014.

In this work, we installed our muon telescope at a distance of 5 km south of the center of the caldera for the purpose of predicting a 2011 Shinmoe-dake eruption. We report our successful VLRM, which exhibits a low-density region below the Shinmoe-dake caldera floor that might cause the caldera to collapse in the future.

2 Principle

2.1 Muography

In this section, we briefly introduce the principle of muography. To measure the internal density distribution of a volcano, the integrated flux of cosmic-ray muons before and after passing through the volcano is compared. Primary cosmic rays, which predominantly consist of protons, arrive at the top of the earth's atmosphere and generate muons as a consequence of interaction with atmospheric nuclei. A number of past experiments have measured the cosmic-ray muon energy spectrum at sea level (e.g. Achard et al., 2004; Haino et al., 2004; Allkofer et al., 1985; Jokisch et al., 1979). Although there are systematic differences between them, the average energy of muons tends to be higher at larger zenith angles. For instance, while the integrated vertical muon flux (0° from zenith) is 7×10^{-3} and $5 \times 10^{-8} \text{ cm}^{-2} \text{ sr}^{-1} \text{ s}^{-1}$ above 1 GeV and 1 TeV, respectively (Beringer et al., 2012), the horizontal flux (85° from zenith) is 1×10^{-4} and $1.6 \times 10^{-7} \text{ cm}^{-2} \text{ sr}^{-1} \text{ s}^{-1}$ above 1 GeV and 1 TeV, respectively (Allkofer et al., 1985). For the purpose of interpolating the measured flux, several groups have formulated approximate equations of the muon energy spectrum as a function of energy and zenith angle (e.g. Gaisser, 2002; Matsuno et al., 1984; Smith and Duller, 1959). However, the systematic difference between them ranges from 10 to 100 % (Lesparre et al., 2010).

The interaction of muons with matter is also well known. The energy loss of muons through matter is given as a function of density-length (density \times path length along the

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



2.2.2 Scattering of decay electrons in the atmosphere

As the distance between the target and muography observation site becomes longer, the atmospheric density-length increases, and thus decay electrons are scattered more often and lose their initial directions. The typical length that the Bremsstrahlung process takes place is called the radiation length (X_0), and X_0 for an electron is 303.9 m in a 1 atm atmosphere. When the distance that a muon has traveled is much longer than its non-relativistic decay length (660 m) and X_0 for an electron, the muon will decay, and its decay electron will experience many Bremsstrahlung processes and will deviate from the original muon's path. Because muons are 207 times heavier than electrons, muons do not experience Bremsstrahlung processes in the atmosphere within this distance scale.

Figure 3 shows the results of extended air shower Monte Carlo (MC) simulations. A MC code called Cosmos (Kasahara and Cohen, 2008) was used to reproduce the energy spectrum of the electromagnetic (EM) components (electron and positron) arriving from the angle region between 50 and 90° at sea level. In these simulations, 9.5×10^8 primary particles with an energy range between 260 MeV and 260 TeV (Beringer et al., 2012) were injected at the top of Earth's atmosphere. The top 5 elements (H, He, C, N, and O) were selected as primary particles. These particles were injected into the atmosphere isotropically within an angle range between 0 and 95° from zenith.

In Fig. 3, the differential flux is plotted within an energy range between 10 MeV and 1 TeV. Electrons with energies above 10 MeV can penetrate a 5 cm thick plastic scintillator, and thus these components could be background sources when thin plastic scintillators (e.g. Lesparre et al., 2012) or gaseous detectors (e.g. Cârloganu et al., 2013; Oláh et al., 2012) are used for muography observations. The EM flux integrated over the energy range above 10 MeV and 1 GeV are 9.4×10^{-5} and $1.7 \times 10^{-6} \text{ cm}^{-2} \text{ sr}^{-1} \text{ s}^{-1}$, respectively. These values are equivalent to the muon flux (arriving at 85° from zenith) after passing through 100 and 1200 m of water, respectively. These electrons and

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

secondary electrons generated inside a lead shield as a function of depth from the surface. 100 electrons with energies of 100 GeV were injected. The number of secondary electrons increased quickly but eventually decreased at a depth of 50 mm from the surface, and they almost disappeared deeper than 450 mm from the surface. 4_{-2}^{+3} electrons remained within a depth region between 400 and 450 mm, and they completely disappeared deep than this region.

We muographically imaged Shinmoe-dake volcano, located in Kagoshima and Miyazaki Prefectures in Japan. The altitudes at our muography telescope and the caldera floor before and after the 2011 eruption were 610, 1240 and 1350 m, respectively (Fig. 6, Shimono et al., 2011). Subsequent to the eruption, no significant changes to the topography of the caldera floor have been reported. We installed our muography telescope 5 km south of the center of the Shinmoe-dake caldera (Fig. 7). Therefore, the elevation angles of the caldera floor before and after the eruption were 128 and 150 mrad, respectively. We defined an azimuth angle of 0 mrad in the direction of the center of the caldera. The angular interval (Tanaka et al., 2014) of the telescope was ± 33 mrad, therefore the telescope could capture the magma deposit beneath the present caldera floor within a 131 ± 33 mrad range of elevation angles. Figure 8 shows the path length distribution of Shinmoe-dake as a function of elevation and azimuth angles. As seen in this figure, path lengths exceeded 5000 m in the region below the 100 mrad elevation angle. It was difficult for us to image this region with our present telescope. The measurements began on 10 October 2014 and continued until 04 November 2014. The observational duration was 64 days.

3.2 Analysis

3.2.1 Muon tracking

The directions of the incident particles in the telescope can be determined by connecting two vertex points in the first and the last position sensitive planes (PSP #1 and PSP

#6). For muon tracking, we selected straight trajectories for incident events according to the following process:

- a. When an event was detected in all 6 PSPs within a time window of 40 ns, the vertex points in PSP#1 and #6 were recorded.
- 5 b. When two or more events were detected in a single PSP within a time window of 40 ns, these events were not recorded (Tanaka et al., 2001).
- c. When the vertex points in the redundant PSPs, i.e. PSPs #2, #3, #4 and #5, deviated within $\pm N$ segments (N value) from the straight line created by connecting the vertex points in (A), event tracks were generated. Smaller N values gave better tracking linearity.

Figure 9 illustrates our tracking scheme for $N = 1$. Figure 10 shows the number of recorded open-sky event tracks that survived for different N values. Because we could not identify the location of the vertex points within the scintillator strip, the telescope failed to track some events if only linear trajectories were selected ($N = 0$). The figure also shows that the number of muons scattered at large angles in the telescope was negligible because there are no significant variations for $N > 2$.

Figure 11 shows the maximum deviations of the Geant4 simulated vertex points in 4 redundant PSPs (i.e. PSP #2, #3, #4, and #5) from a linear trajectory drawn by connecting the vertex points in PSPs #1 and #6. For these simulations, muons were injected into the virtual telescope, which is identical to the actual telescope, with an energy spectrum of 70° from zenith. The plot in Fig. 11 is consistent with the N -dependence of the recorded number of tracks (Fig. 10), and we therefore conclude that the event tracks recorded with our telescope using the just discussed tracking scheme were muon tracks. We specified $N = 1$, because large N values blur the resultant image.

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and H. K. M. Tanaka

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



at high elevations, the scattering of backward-directed muons is negligible. However, for LRM, because the region of interest is usually located at low elevations, this effect also becomes serious.

To estimate muon scatterings inside the telescope, MC simulations (Geant4) were performed. In these simulations, 500 000 muons were injected into the modeled radiation shields (50 cm thick lead and 30 cm thick stainless steel plates) according to an energy spectrum measured within an angle range between 60 and 70° from zenith (Achard et al., 2004; Tsuji et al., 1998; Abdel-Monem et al., 1975). Figure 12 shows the scattering angle distribution of near-horizontal muons after they passed through the radiation shields. While approximately 4% of the muons scattered at angles greater than 33 mrad, only 0.7% of the muons scattered at angles greater than 100 mrad. These numbers respectively correspond to scattered muon fluxes of 4×10^{-6} and $7 \times 10^{-7} \text{ cm}^{-2} \text{ sr}^{-1} \text{ s}^{-1}$ if we assume that the arriving angle of the muons was 80°.

To estimate background level from muon scattering inside the telescope, we calculated the flux of muons that scattered towards the telescope's y direction. The elevation angular dependence of the number of background events $N_{\text{BG}}(\theta)$ measured within the elevation angle range θ to $\theta + \Delta\theta$ is

$$N_{\text{BG}}(\theta) = \int N(\theta)P(\theta)d\theta, \quad (2)$$

where $\Delta\theta$ is an angular interval of our detector, and $N(\theta)$ is the elevation angular dependence of the number of muons measured within an elevation angle range θ to $\theta + \Delta\theta$. $P(\theta)$ is given by

$$P(\theta) = \frac{\int_{\theta}^{\theta+\Delta\theta} \phi(\theta)d\theta}{\int_0^{\infty} \phi(\theta)d\theta}, \quad (3)$$

where $\phi(\theta)$ is the muon scattering probability distribution function shown in Fig. 12.

The elevation angular dependence of the forward-directed and backward-directed background event flux is plotted in Fig. 13 by calculating Eq. (2). Because low energy

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



than at Satsuma-Iwojima or (B) background events offset the data. The path lengths of the muons at Shinmoe-dake ranged from 5000 to 7000 m within the angular region. The corresponding flux was $\sim 10^{-10} \text{ cm}^{-2} \text{ sr}^{-1} \text{ s}^{-1}$, therefore (A) can be rejected. Because the ridge of Shinmoe-dake is located at an elevation angle of approximately only 200 mrad, even small muon scatterings inside the telescope could appear to be more horizontal muon events from the direction below the ridge. Therefore, the possible background sources were the sum of the (a) forward-directed and (b) backward-directed muon scatterings inside the detector. Figure 15 compares the elevation angular dependence of the Shinmoe-dake flux with the sum of the forward- and backward-directed background event fluxes $N_{\text{BG}}(\theta)$ for $N = 1$. As seen in this figure, the excess of the Shinmoe-dake flux at lower elevation angles (33 and 66 mrad) can be explained by these backgrounds. The difference between the Shinmoe-dake flux and $N_{\text{BG}}(\theta)$ at elevation angles larger than 100 mrad was recognized as muons after passing through the volcano.

4.2 Very long range muographic imaging of the Shinmoe-dake volcano

To derive the average density along the muon paths passing through a region right below the caldera floor, in Fig. 16 we compared the theoretical and experimental Shinmoe-dake muon fluxes at an elevation angle of 130 mrad as a function of azimuth angle (ϕ). In this figure, only the data points with more than 50 events were plotted. For $\phi < -350$ mrad and $\phi > 200$ mrad, the numbers of events were less than 50 because (a) the active areas of the telescope were small, and (b) the rock became thick.

5 Discussion

Figure 17 shows muographic images below the Shinmoe-dake caldera. These images were produced by interpolating the discrete data points for the purpose of enhancing the visual effect. In this image, because we did not record sufficient number of events

GID

5, 207–246, 2015

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



below the caldera floor before the 2011 eruption, we modeled the density in this region by applying the average density of the crater rim. In Fig. 17, an east–west cross section in the region of the mountain peak is also plotted to show the profile of the caldera before and after the 2011 eruption. To illustrate the statistical significance of the density contrast, the 1σ , 2σ and 3σ upper limits of the density are plotted in Fig. 17b–d, respectively.

As seen in Fig. 17, the density is sensitive to variations of the sigma values at an elevation angle of 100 mrad. This means that the statistic error is large; therefore there is a large uncertainty in the density values in this region.

As seen in Fig. 17, there are two low-density regions below the present caldera floor. The regions are localized at the eastern and western areas, while the central area has higher densities. Figure 18 shows the location of a crater of an explosive eruption (subplinian event) and a growing lava dome found on the caldera floor on the morning of 28 January 2011 (Nakada et al., 2013). A series of eruptions in 2011 at Shinmoe-dake started from this crater.

Figure 19 shows the results of airborne synthetic aperture radar (SAR) measurements in February and October 2011 (Kobayashi et al., 2013). Figure 19 shows the vertical cross sections along the lines shown in Fig. 18. As seen in this figure, the southeast part of the caldera floor collapsed some time between February and October 2011. The size of the collapse was 100 m in diameter and 30 m in depth; however, the depth may have been greater. The location of the collapse coincides with that of the steam vent shown in Fig. 18.

In the present analysis, the caldera topography measured in February 2011 was utilized to derive the average density along the muon paths. This means that the 100 m diameter collapse was not incorporated into our analysis, and the muographic images shown in Fig. 17 would reflect this newly created vertical channel. When considering that the typical path length in this region is ~ 1000 m, the reduction in density would be $\sim 10\%$. This reduction could explain the low-density region found in the eastern part of the caldera. However, there were large density errors in this region, therefore the

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

contrast was not as apparent (the density contrast disappears when we take the 3 σ upper limit of the determined density; Fig. 17d).

Unlike the eastern part, the topography of the western part of the caldera floor seems more stable and did not change between February and October 2011 (Fig. 19). However, muographic images show a distinct low-density region immediately below the western caldera floor (Fig. 17). As a consequence, the following scenario can explain the sequence of the 2011 Shinmoe-dake eruption. Lava and ash emitted from the Shinmoe-dake caldera almost filled it over less than a month after the first eruption on 19 January 2011, which took place in the western part of the caldera. After the volcano became less active at the end of February, the more fragile part inside the caldera (i.e. the layer of thickly deposited ash shown in Fig. 18) collapsed right above a steam vent (Fig. 18) and created a 100 m diameter vertical channel. On the other hand, a lava layer covered an initial subplinian event crater (Fig. 18), and hotter and more fluid magma underneath the solidified surface (that is more stable than deposited ash) was gradually drained into the crater. As a consequence, a low-density (porous) region was created underneath the western caldera floor.

In prior works, it was reported that a void was created beneath the caldera floor of Miyake-jima volcano in 2000. The day after the 8 June eruption, an ellipsoidal region with a length of ~ 700 m and width of ~ 400 m collapsed ~ 200 m in depth (Nakada et al., 2001). The precursor of the caldera collapse was detected by gravimetric measurements 2 days prior to the event. Furuya et al. (2001) compared the spatio-temporal changes between 2 days and 2 years before the collapse. A difference in gravity values between those times was $-145 \mu\text{gal}$ inside the caldera, and the data points showed that the values below $-100 \mu\text{gal}$ were located within a region 1.0–1.5 km from the caldera center. A gravity change of $-145 \mu\text{gal}$ corresponds to an upheaval of several tens of centimeters. However, such an upheaval was not detected using global positioning system (GPS) measurements. Furuya et al. (2001) interpreted this inconsistency as a new void that formed underneath the crater floor during the 2000 Miyake-jima eruption.

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In the present work, muography measurements were carried out at a distance of 5 km from the volcano summit. This implies that VLRM can provide us with a new tool to predict caldera collapse by mapping out the density distribution directly beneath the caldera floor. On 8 October 2014, it was reported that Shinmoe-dake began to deform because of an accumulation of magma beneath the volcano (Geospatial Information Authority of Japan, 2014) and reached the same level of expansion that was observed right before the 2011 Shinmoe-dake eruption. In the future, joint measurements between muographic and absolute gravimetric measurements can provide more useful information for predicting volcano eruption sequences by independently measuring magma movements in shallow regions compared to movements in deeper regions. Such measurements can give us new information about magma dynamics inside volcanoes.

6 Conclusions

In this work, we successfully applied a calorimeter-type muography telescope with radiation shields and redundant detectors (PSPs) to an erupting volcano for very long-range muography (VLRM). Conventional muography is limited to the vicinity of a volcanic crater, and it is therefore impractical to apply conventional techniques when monitoring erupting volcanoes. Our results show that muography has the potential to be a practical tool for monitoring hazardous volcanic eruptions.

Entry to the summit area of Shinmoe-dake volcano is still restricted. Since the beginning of 2011 eruption, the surface phenomena associated with eruption transitions have been revealed by airborne and land-based measurements. However, it has been technically difficult to image the internal structure of the volcano from a secure distance. Using VLRM, we found a low-density region directly below the caldera floor, which implies that magma dynamics occurred below the caldera floor after eruptions became less active in February 2011. The location of the low-density region is coincident with

the crater that was formed during the initial eruption in January 2011. The void might cause a caldera collapse.

Our present proof-of-principle (POP) work shows the capability of conducting muographic measurements 5 km from the summit. This is considered to be a safe distance for many volcanic eruptions. On the other hand, for muography applied to erupting volcanoes, precise topographic information of the target is required because the surface topography of an erupting volcano varies with time. Such information can be obtained using repeated airborne synthetic aperture radar (SAR) measurements. In conjunction with recent progress in those airborne techniques, the range of application of muography would be remarkably expanded.

Acknowledgements. We thank Kirishima Heights for offering the observation site and helping the muographic observation.

References

- Abdel-Monem, M. S., Benbrook, J. R., Osborne, A. R., Sheldon, W. R., Choate, L. M., Magnuson, C. E., Duller, N. M., and Green, P. J.: Cosmic ray muon spectra at zenith angles 65° and 80° using the AMH Magnetic Spectrometer, in: vol. 6, 14th International Cosmic Ray Conference, München, p. 2043, 1975.
- Achard, P., Adriani, O., Aguilar-Benitez, M. et al.: Measurement of the atmospheric muon spectrum from 20 to 3000 GeV, Phys. Lett. B, 598, 15–32, doi:10.1016/j.physletb.2004.08.003, 2004.
- Agostinelli, S., Allison, J., Amako, K. et al.: Geant4 – a simulation toolkit, Nucl. Instrum. Meth. A, 506, 250–303, doi:10.1016/S0168-9002(03)01368-8, 2003.
- Allkofer, O. C., Bella, G., Dau, W. D., Jokisch, H., Klemke, G., Oren, Y., and Uhr, R.: Cosmic ray muon spectra at sea-level up to 10 TeV, Nucl. Phys. B, 259, 1–18, doi:10.1016/0550-3213(85)90294-9, 1985.
- Barrett, P. H., Bollinger, L. M., Cocconi, G., Eisenberg, Y., and Greisen, K.: Interpretation of cosmic-ray measurements far underground, Rev. Mod. Phys., 24, 133–178, 1952.
- Beringer, J., Arguin, J.-F., Barnett, R. M. et al.: Review of particle physics, Phys. Rev. D, 86, 010001, doi:10.1103/PhysRevD.86.010001, 2012.

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Carbone, D., Gibert, D., Marteau, J., Diament, M., Zuccarello, L., and Galichet, E.: An experiment of muon radiography at Mt Etna (Italy), *Geophys. J. Int.*, 196, 633–643, doi:10.1093/gji/ggt403, 2013.

Cârloganu, C., Niess, V., Béné, S., Busato, E., Dupieux, P., Fehr, F., Gay, P., Miallier, D., Vulpescu, B., Boivin, P., Combaret, C., Labazuy, P., Laktineh, I., Lénat, J.-F., Mirabito, L., and Portal, A.: Towards a muon radiography of the Puy de Dôme, *Geosci. Instrum. Method. Data Syst.*, 2, 55–60, doi:10.5194/gi-2-55-2013, 2013.

De Lellis, G., Consiglio, L., Strolin, P. et al.: Muon radiography with nuclear emulsions: stromboli and other projects, in: MUOGRAPHERS2014, Tokyo, 2014.

Furuya, M., Okubo, S., Tanaka, Y., Sun, W., Watanabe, H., Oikawa, J., and Maekawa, T.: Caldera formation process during the Miyakejima 2000 volcanic activity detected by spatio-temporal gravity changes, *J. Geography*, 110, 217–225, 2001.

Gaisser, T. K.: Semi-analytic approximations for production of atmospheric muons and neutrinos, *Astropart. Phys.*, 16, 285–294, doi:10.1016/S0927-6505(01)00143-8, 2002.

Geospatial Information Authority of Japan: Press release on 08 October, see <http://www.gsi.go.jp/WNEW/PRESS-RELEASE/2014-goudou1008.html>, last access: 16 December 2014.

Groom, D. E., Mokhov, N. V., and Striganov, S. I.: Muon stopping power and range tables 10 MeV–100 TeV, *Atom Data Nucl. Data*, 78, 183–356, doi:10.1006/adnd.2001.0861, 2001.

Haino, S., Sanuki, T., Abe, K., Anraku, K., Asaoka, Y., Fuke, H., Imori, M., Itasaki, A., Maeno, T., Makida, Y., Matsuda, S., Matsui, N., Matsumoto, H., Mitchell, J. W., Moiseev, A. A., Nishimura, J., Nozaki, M., Orito, S., Ormes, J. F., Sasaki, M., Seo, E. S., Shikaze, Y., Stremtner, R. E., Suzuki, J., Takasugi, Y., Tanaka, K., Tanizaki, K., Yamagami, T., Yamamoto, A., Yamamoto, Y., Yamato, K., Yoshida, T., and Yoshimura, K.: Measurements of primary and atmospheric cosmic-ray spectra with the BESS-TeV spectrometer, *Phys. Lett. B*, 594, 35–46, doi:10.1016/j.physletb.2004.05.019, 2004.

Hernández, I., Hernández, P., Pérez, N., Tanaka, H., Miyamoto, S., Barrancos, J., and Padrón, E.: Application of emulsion imaging system for cosmic-ray muon radiography to explore the internal structure of Teide and Cumbre Vieja volcanoes in the Canary Islands, Spain, in: EGU General Assembly 2013, Vienna, available at: <http://adsabs.harvard.edu/abs/2013EGUGA..15..953H> (last access: 14 January 2015), 2013.

Japan Meteorological Agency: Press release on 03 February, available at: <http://www.jma.go.jp/jma/press/1102/03a/yochiren110203.html>, available at: <http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:No+Title#0> (last access: 15 December 2014), 2013.

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[▶⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Jokisch, H., Carstensen, K., Dau, W., Meyer, H., and Allkofer, O.: Cosmic-ray muon spectrum up to 1 TeV at 75° zenith angle, *Phys. Rev. D*, 19, 1368–1372, doi:10.1103/PhysRevD.19.1368, 1979.

Kasahara, K. and Cohen, F.: A new parallel processing scheme enabling full Monte Carlo EAS simulation in the GZK Energy Region, in: 30th International Cosmic Ray Conference, vol. 4, 581–584, available at: <http://adsabs.harvard.edu/abs/2008ICRC....4..581K> (last access: 15 December 2014), 2008.

Kobayashi, T., Umehara, T., Uemoto, J., and Satake, M.: Volcanic monitoring by polarimetric and interferometric airborne SAR (Pi-SAR2), in: 2013 Asia-Pacific Conference on Synthetic Aperture Radar (AP SAR), Tsukuba, 487–490, 2013.

Kusagaya, T., Tanaka, H., Taketa, A., Maekawa, T., and Yokoyama, I.: Development of a multifold segmented muon detection system to improve the maximum resolvable distance of muography, in: Japan Geoscience Union Meeting 2013, Chiba, 2013.

Lesparre, N., Gibert, D., Marteau, J., Déclais, Y., Carbone, D., and Galichet, E.: Geophysical muon imaging: feasibility and limits, *Geophys. J. Int.*, 183, 1348–1361, doi:10.1111/j.1365-246X.2010.04790.x, 2010.

Lesparre, N., Gibert, D., Marteau, J., Komorowski, J.-C., Nicollin, F., and Coutant, O.: Density muon radiography of La Soufrière of Guadeloupe volcano: comparison with geological, electrical resistivity and gravity data, *Geophys. J. Int.*, 190, 1008–1019, doi:10.1111/j.1365-246X.2012.05546.x, 2012.

Matsuno, S., Kajino, F., Kawashima, Y., Kitamura, T., Mitsui, K., Muraki, Y., Ohashi, Y., Okada, A., Suda, T., Minorikawa, Y., Kobayakawa, K., Kamiya, Y., Nakamura, I., and Takahashi, T.: Cosmic-ray muon spectrum up to 20 TeV at 89° zenith angle, *Phys. Rev. D*, 29, 1–23, doi:10.1103/PhysRevD.29.1, 1984.

Miyamoto, S., Bozza, C., Ambrosio, N. D., De Lellis, G., Nakamura, M., Rescigno, R., Russo, A., Shimizu, H., Sirignano, C., Stellacci, S., Strolin, P., Tanaka, H., and Tioukov, V.: The first radiography result of the latest lava dome in Unzen by cosmic muons, in: EGU General Assembly 2012, Vienna, available at: <http://adsabs.harvard.edu/abs/2012EGUGA..14.1938M> (last access: 14 January 2015), 2012.

Nakada, S., Nagai, M., Yasuda, A., Shimano, T., Geshi, N., Ohno, M., Akimasa, T., Kaneko, T., and Fujii, T.: Chronology of the Miyakejima 2000 eruption: characteristics of summit collapsed crater and eruption products, *J. Geog.*, 110, 168–180, 2001.

**Technique to explore
the internal structure
of an erupting
volcano**T. Kusagaya and
H. K. M. Tanaka

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nakada, S., Nagai, M., Kaneko, T., Suzuki, Y., and Maeno, F.: The outline of the 2011 eruption at Shinmoe-dake (Kirishima), Japan, *Earth Planets Space*, 65, 475–488, doi:10.5047/eps.2013.03.016, 2013.

Oláh, L., Barnaföldi, G. G., Hamar, G., Melegh, H. G., Surányi, G., and Varga, D.: CCC-based muon telescope for examination of natural caves, *Geosci. Instrum. Method. Data Syst.*, 1, 229–234, doi:10.5194/gi-1-229-2012, 2012.

Shimono, T., Minami, H., Nishii, Y., Ohno, H., and Watanabe, K.: Observation of volcanic crater of Mt. Kirishima (Shinmoedake) by Airborne SAR, *Bull. Geospatial Inf. Auth. Japan*, 121, 189–194, 2011.

Smith, J. A. and Duller, N. M.: Effects of pi meson decay-absorption phenomena on the high-energy mu meson zenithal variation near sea level, *J. Geophys. Res.*, 64, 2297, doi:10.1029/JZ064i012p02297, 1959.

Tanaka, H. K. M.: Particle geophysics, *Annu. Rev. Earth Pl. Sc.*, 42, 535–549, doi:10.1146/annurev-earth-060313-054632, 2014.

Tanaka, H. K. M. and Yokoyama, I.: Possible application of compact electronics for multilayer muon high-speed radiography to volcanic cones, *Geosci. Instrum. Method. Data Syst.*, 2, 263–273, doi:10.5194/gi-2-263-2013, 2013.

Tanaka, H. K. M., Nagamine, K., Kawamura, N., Nakamura, S. N., Ishida, K., and Shimomura, K.: Development of the cosmic-ray muon detection system for probing internal-structure of a volcano, *Hyperfine Interact.*, 138, 521–526, doi:10.1023/A:1020843100008, 2001.

Tanaka, H. K. M., Nakano, T., Takahashi, S., Yoshida, J., Takeo, M., Oikawa, J., Ohminato, T., Aoki, Y., Koyama, E., Tsuji, H., and Niwa, K.: High resolution imaging in the inhomogeneous crust with cosmic-ray muon radiography: the density structure below the volcanic crater floor of Mt. Asama, Japan, *Earth Planet. Sc. Lett.*, 263, 104–113, doi:10.1016/j.epsl.2007.09.001, 2007a.

Tanaka, H. K. M., Nakano, T., Takahashi, S., Yoshida, J., Ohshima, H., Maekawa, T., Watanabe, H., and Niwa, K.: Imaging the conduit size of the dome with cosmic-ray muons: the structure beneath Showa-Shinzan Lava Dome, Japan, *Geophys. Res. Lett.*, 34, L22311, doi:10.1029/2007GL031389, 2007b.

Tanaka, H. K. M., Uchida, T., Tanaka, M., Shinohara, H., and Taira, H.: Cosmic-ray muon imaging of magma in a conduit: degassing process of Satsuma-Iwojima Volcano, Japan, *Geophys. Res. Lett.*, 36, 1–5, doi:10.1029/2008GL036451, 2009a.

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[▶⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Tanaka, H. K. M., Uchida, T., Tanaka, M., Takeo, M., Oikawa, J., Ohminato, T., Aoki, Y., Koyama, E., and Tsuji, H.: Detecting a mass change inside a volcano by cosmic-ray muon radiography (muography): first results from measurements at Asama volcano, Japan, *Geophys. Res. Lett.*, 36, L17302, doi:10.1029/2009GL039448, 2009b.

5 Tanaka, H. K. M., Taira, H., Uchida, T., Tanaka, M., Takeo, M., Ohminato, T., Aoki, Y., Nishiyama, R., Shoji, D., and Tsuji, H.: Three-dimensional computational axial tomography scan of a volcano with cosmic ray muon radiography, *J. Geophys. Res.*, 115, B12332, doi:10.1029/2010JB007677, 2010.

10 Tanaka, H. K. M., Kusagaya, T., and Shinohara, H.: Radiographic visualization of magma dynamics in an erupting volcano, *Nat. Commun.*, 5, 3381, doi:10.1038/ncomms4381, 2014.

Tsuji, S., Katayama, T., Okei, K., Wada, T., Yamamoto, I., and Yamashita, Y.: Measurements of muons at sea level, *J. Phys. G Nucl. Partic.*, 24, 1805–1822, doi:10.1088/0954-3899/24/9/013, 1998.

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka




Figure 1. Visual comparison with prior works. The past muography observation sites are plotted as distance from the peak of the volcano.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka




Figure 2. Elevation angular dependence of muon flux after passing through a given rock thickness. The rock thickness is given in units of meter water equivalent (m.w.e.).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Technique to explore
the internal structure
of an erupting
volcano**T. Kusagaya and
H. K. M. Tanaka


[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Figure 3. Differential energy spectrum of electromagnetic components (electron and positron) of secondary cosmic rays at sea level. The arriving angles range between 50 and 90° from zenith.

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka




Figure 4. Schematic of the muography telescope used for the present work.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka




Figure 5. Stopping distribution of electrons inside a lead block. The incident energy is 100 GeV. The stopping probability per 1 mm at different depths (red) and the corresponding cumulative probability (blue) are shown.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka




Figure 6. Caldera deformation captured with airborne synthetic aperture radar (SAR) measurements. The black dotted and blue solid lines respectively show the topography of the caldera floor before (March 2009) and after (February 2011) the 2011 eruption for the west–east cross section **(a)** and south–north cross section **(b)**.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




Figure 7. Geometrical configuration of the experimental setup. The left panel shows the location of the Shinmoe-dake volcano. The right panel shows an enlarged map in the vicinity of the Shinmoe-dake volcano. The muography telescope (Mu) was located 5 km south southwest from the center of the Shinmoe-dake volcano caldera. The topographic map does not reflect the land deformation during the 2011 eruption.

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka




Figure 8. Path length distribution of volcanoes viewed from different distances. The solid angle when viewing a larger volcano (Shinmoe-dake) **(a)** is smaller than that when viewing a smaller volcano (Satsuma-Iwojima) **(b)** if the distance to the target is farther.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka




Figure 9. Example of muon tracking for $N = 1$. If a signal (a red square) is generated at $(x, y) = (5, 10)$ in PSP #1 (a) and at $(x, y) = (15, 1)$ in PSP #6 (f) simultaneously, the signal generation points (red squares) in PSPs #2–5 (b–e) are used for tracking.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Technique to explore
the internal structure
of an erupting
volcano**T. Kusagaya and
H. K. M. Tanaka


[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Figure 10. Number of event tracks as a function of N . Error bars indicate the range of integration.

**Technique to explore
the internal structure
of an erupting
volcano**T. Kusagaya and
H. K. M. Tanaka


[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Figure 11. Muon scattering inside the telescope. The event was normalized to the total number of events.

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka




Figure 12. Scattering angle distribution of near-horizontal muons. The events are normalized by the total number of muons that were injected.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




Figure 13. Elevation angular dependence of background event flux. The background levels are expected to be measured for different elevation angles with the telescope and are plotted for the events arriving from the direction of the target mountain **(a)** and from the side opposite the mountain **(b)**. The plot also shows the background event flux before (red squares) and after (blue diamonds) selecting linear trajectories from these events with a selection criterion of $N = 1$.

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka




Figure 14. Elevation angular dependence of averaged event flux observed at Shinmoe-dake and Satsuma-Iwojima. The flux was averaged between azimuth ranges of -430 to 430 mrad for Shinmoe-dake and 0 to 400 mrad for Satsuma-Iwojima. The backward-directed (open sky) and forward-directed (mountain side) flux is shown for Shinmoe-dake and Satsuma-Iwojima with blue and red triangles and blue and red circles, respectively.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka




Figure 15. Comparison of the measured and theoretical background fluxes. The blue and red circles shows the angular dependence of the Shinmoe-dake and Iwojima fluxes, and the purple crosses and green circles show the estimated background from backward-directed muons and both forward and backward-directed muons, respectively.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka




Figure 16. Azimuth distribution of event flux measured at the Shinmoe-dake volcano. The data collected at elevation angle of 130 mrad are plotted along with the theoretical flux calculated for different density values (1.5, 2.0, 2.5, and 3.0 g cm^{-3}). The data before (black squares) and after (red squares) the background event removal are plotted.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 17. Interpolated muographic images showing (a), mean densities, (b), 1σ , (c), 2σ , and (d), 3σ upper limit densities below the Shinmoe-dake caldera floor. The east–west cross sections of the mountain before (dotted lines) and after the 2011 eruption (February 2011) are shown. The color scale shows the average density along the muon paths.

Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka




Figure 18. Schematic of the Shinmoe-dake caldera after the 2011 eruption (Nakada et al., 2013). The left and right panels respectively show the caldera immediately after the eruption (28 January 2011) and after more than a year had passed (the summer of 2012). The numbers in the plot indicate: 1, a vent of subplinian events, 2, a vent of continuous emission of fine ash, 3, a lava dome, 4, a steam vent, 5, the limit of lava accumulation, and 6, altered, thickly deposited ash. The vertical cross sections are plotted along 3 red horizontal lines (a–c) in Fig. 19

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Technique to explore the internal structure of an erupting volcano

T. Kusagaya and
H. K. M. Tanaka

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




Figure 19. Vertical cross sections inside the caldera of Shinmoe-dake volcano along the lines (a–c) in Fig. 18 on 6 October 2011 (red) and 26 February 2011 (blue) by airborne SAR (Pi-SAR2) measurements (Kobayashi et al., 2013).